



#8

Form P10-1449			
ATTY DOCKET NO. 32-95	SERIAL NO. 08/477,354	FILING DATE June 7, 1995	
APPLICANT Hawley-Nelson et al.	·	GROUP Unassigned	

U.S. PATENT DOCUMENTS

Examinr. Initial	Document Number	Date	Name	Class	Subclass	Filing Date if Appropriate
25	04,946,787	Aug 7, 1990	Eppstein et al.			,
IBB	08/069,720	June 1, 1993	Jessee et al.			1
183	08/090,290	July 12, 1993	Jessee et al.			
BB	08/274,397	July 12, 1994	Jessee et al.			

FOREIGN PATENT DOCUMENTS

		Document Number	Date	Country	Class	Subclass	Translation Yes/No
P.C.	./	WO A91/16024	31.10.91				
MB	1	WO 93/07282	04.15.93				
BB		WO 93/07283	04.15.93				

OTHER PRIOR ART (including Author, Title, Date, Pertinent Pages, etc.)

B		/	DeRoberts et al., "Intracellular migration of nuclear proteins in Xenopus oocytes," Nature <u>272</u> :254-256 (1978).
P B			Väänänen et al., "Fusion and Haemolysis of Erythrocytes Caused by Three -Togaviruses: Semiki Forest, Sindbis, and Rubella," J. Gen. Virology (1980), <u>46</u> : 467-475.
JB		ر بہ	Carrasco, L. et al. "Modification of Membrane Permeability in Vaccinia Virus- Infected Cells," (1982), J. Virol. <u>117</u> :62-69.
183		/	Eytan, G.D., "Use of Liposomes for Reconstitution of Biological Functions," Biochem. Biphys. Acata (1982) <u>694</u> :185-202.
J&D		_	Young et al., "Interaction of Enveloped Viruses with Planar Bilayer Membranes: Observations on Sendai, Influenza, Vesicular Stomatitis, and Simiki Forest Viruses," Virology (1983) 128:186-194.
SB			Marsh et al., "Interactions of Simiki Forest Virus Spike Glycoprotein Rosettes and Vesicles with Cultured Cells," J. Cell Biol. (1983) <u>96</u> :455-461.
gB		1	Schlegel, R. et al. "Inhibition of VSV Binding and Infectivity by Phosphatidylserine: Is Phosphatidylserine a VSV-Binding Site?" Cell 32:639-646 (1983).
8-13		_	Kalderon et al., "A Short Amino Acid Sequence Able to Specify Nuclear Location," Cell <u>39</u> :499-509 (1984).
JES .		1	Kraaijeveld, S.A. et al., "The effect of liposomal charge on the neutralizing antibody response against inactivated encephalomyocarditis and Simiki Forest Viruses," Clin. Exp. Immunol., (1984)56:509-514.
RB	1	/	Schlegel, R. and M. Wade, "Biologically Active Peptides of the Vesicular Stomatitus Virus Glycoprotein," J. Virol. <u>53</u> (1):319-323 (1985).
SB			Klappe, K. et al., "Parameters Affecting Fusion between Sendai Virus and Liposomes. Role of Virul Proteins, Liposome Composition, and pH," Biochemistry (1986) <u>25</u> :8252-8260.
BB		1	Sands, J.A., "Virucidal activity of cetyltrimethylammonium bromide below the critical micelle concentration," FEMS Microbiol. Lett. (1986) 36:261-263.
JB			Scheule, "Novel Preparation of Functional Sindbis Virosomes," Biochemistry (1986) 25:4223-4232.
		7	

EXAMINER (

DATE CONSIDERED 4/84

*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

ATTY DOCKET NO. 32-95 SERIAL NO. 08/477,354 FILING DATE June 7, 1995 APRI ICANIT. Hawley-Nelson et al. GROUP Unassigned	MAST ROOM		Sheet <u>2</u> of <u>3</u>
ATTY DOCKET NO. 32-95 SERIAL NO. 08/47/,334 FILLING DATE Julie 1, 334	TOWN TO THE STATE OF THE STATE	ETI INC DATE June 7, 1995	
	ATTY DOCKET NO. 32-95 SERIAL NO. 08/477,334 APPLICANT Hawley-Nelson et al.		

OTHER PRIOR ART (including Author, Title, Date, Pertinent Pages, etc.)

Biotech. 2:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098.	OTHER PRIOR ART (including Author, Title, Date, Pertinent Pages, etc.)				
Macromolecules into Cells Using HVI (Sendai virus) Liposomes with Gangliosides, "Exp. Cell Res. (1987) 173:56-69. Otero, M.J., and Carrasco, L. "Proteins are Cointernalized with Virion Particles during Early Infection," (1987), J. Virol. 160:75-80. Tikchonenko, T., et al., (1988) "Transfer of condensed viral DNA into eukaryotic cells using proteoliposomes," Gene 63:321-330. Gould-Fogerite, S. et al., "Chimerasome-mediated gene transfer in vitro and in vivo," (1989) Gene 84:429-438. Kaneda et al., "Introduction and Expression of the Human Insulin Gene in Adult Rat Liver," J. Biol. Chem. (1989) 264(21):1216-1219. Neugebauer, J. "Detergents: An Overview," Meth.Enzymol., (1990) 182:239-253. Lapidot et al., "Fusion-Mediated Microinjection of Liposome-Enclosed DNA into Cultured Cells with the Aid of Influenza Virus Glycoproteins," Experimental Cell Research (1990) 189:241-246. Konopka, K. et al., "Enhancement of human immunodeficiency virus type 1 infection by cationic liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. (1991) 72:2685-2696. Curiel, D.T. et al., "Adenovirus enhancement of transferrin-polylysine-mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8850-8854. Liljistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semiliki Forest Virus Replicon," (1991) Biotech. 2:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Poly	JB.			Peptide Homologous to the SV40 T Antigen Transport Signal, Cell 46:575-582 (1986).	
Particles during Early Infection," (1987), J. Virol. 1891:3-80. Tikchonenko, T., et al., (1988) "Transfer of condensed viral DNA into eukaryotic cells using proteoliposomes," Gene 63:321-330. Gould-Fogerite, S. et al., "Chimerasome-mediated gene transfer in vitro and in vivo," (1989) Gene 84:429-438. Kaneda et al., "Introduction and Expression of the Human Insulin Gene in Adult Rat Liver," J. Biol. Chem. (1989) 264(21):1216-1219. Neugebauer, J. "Detergents: An Overview," Meth.Enzymol., (1990) 182:239-253. Lapidot et al., "Fusion-Mediated Microinjection of Liposome-Enclosed DNA into Cultured Cells with the Aid of Influenza Virus Glycoproteins," Experimental Cell Research (1990) 189:241-246. Konopka, K. et al., "Enhancement of human immunodeficiency virus type 1 infection by cationic-liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. (1991) 72:2685-2696. Curiel, D.T. et al. "Adenovirus enhancement of transferrin-polylysine-mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 82:8850-8854. Liljistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semliki Forest Virus Replicon," (1991) Biotech. 9:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	80			Macromolecules into Cells Using HVJ (Sendai virus) Liposomes with	
eukaryotic cells using proteoliposomes," Gene 63:321-330. Gould-Fogerite, S. et al., "Chimerasome-mediated gene transfer in vitro and in vivo," (1989) Gene 84:429-438. Kaneda et al., "Introduction and Expression of the Human Insulin Gene in Adult Rat Liver," J. Biol. Chem. (1989) 264(21):1216-1219. Neugebauer, J. "Detergents: An Overview," Meth.Enzymol., (1990) 182:239-253. Lapidot et al., "Fusion-Mediated Microinjection of Liposome-Enclosed DNA into Cultured Cells with the Aid of Influenza Virus Glycoproteins," Experimental Cell Research (1990) 189:241-246. Konopka, K. et al., "Enhancement of human immunodeficiency virus type 1 infection by cationic liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. (1991) 72:2685-2696. Curiel, D.T. et al. "Adenovirus enhancement of transferrin-polylysine-mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8850-8854. Liljistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semliki Forest Virus Replicon," (1991) Biotech. 2:1336-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	bs.			Otero, M.J., and Carrasco, L. "Proteins are Cointernalized with Virion Particles during Early Infection," (1987), J. Virol. <u>160</u> :75-80.	
vitro and in vivo," (1989) Gene 84:429-438. Kaneda et al., "Introduction and Expression of the Human Insulin Gene in Adult Rat Liver," J. Biol. Chem. (1989) 264(21):1216-1219. Neugebauer, J. "Detergents: An Overview," Meth. Enzymol., (1990) 182:239-233. Lapidot et al., "Fusion-Mediated Microinjection of Liposome-Enclosed DNA into Cultured Cells with the Aid of Influenza Virus Glycoproteins," Experimental Cell Research (1990) 189:241-246. Kkonopka, K. et al., "Enhancement of human immunodeficiency virus type 1 infection by cationic liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. (1991) 72:2685-2696. Curiel, D.T. et al. "Adenovirus enhancement of transferrin-polylysine-mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8850-8854. Liljistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semliki Forest Virus Replicon," (1991, Biotech. 9:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	12/5			eukaryotic cells using proteoliposomes," Gene <u>65</u> :321-330.	
in Adult Rat Liver," J. Biol. Chem. (1989) 264(21):1716-1719. Neugebauer, J. "Detergents: An Overview," Meth.Enzymol., (1990) 182:239-253. Lapidot et al., "Fusion-Mediated Microinjection of Liposome-Enclosed DNA into Cultured Cells with the Aid of Influenza Virus Glycoproteins," Experimental Cell Research (1990) 189:241-246. Konopka, K. et al., "Enhancement of human immunodeficiency virus type 1 infection by cationic liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. (1991) 72:2685-2696. Curiel, D.T. et al. "Adenovirus enhancement of transferrin-polylysine-mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8850-8854. Litjistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semliki Forest Virus Replicon," (1991) Biotech. 9:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	1265			Gould-Fogerite, S. et al., "Chimerasome-mediated gene transfer in vitro and in vivo," (1989) Gene <u>84</u> :429-438.	
Lapidot et al., "Fusion-Mediated Microinjection of Liposome-Enclosed DNA into Cultured Cells with the Aid of Influenza Virus Glycoproteins," Experimental Cell Research (1990) 189:241-246. Konopka, K. et al., "Enhancement of human immunodeficiency virus type 1 infection by cationic liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. (1991) 72:2685-2696. Curiel, D.T. et al. "Adenovirus enhancement of transferrin-polylysine-mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8850-8854. Liljistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semliki Forest Virus Replicon," (1991) Biotech. 9:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	23			Kaneda et al., "Introduction and Expression of the Human Insulin Gene in Adult Rat Liver," J. Biol. Chem. (1989) <u>264(21)</u> :1216-1219.	
DNA into Cultured Cells with the Aid of Influenza Virus Glycoproteins," Experimental Cell Research (1990) 189:241-246. Konopka, K. et al., "Enhancement of human immunodeficiency virus type 1 infection by cationic-liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. (1991) 72:2685-2696. Curiel, D.T. et al. "Adenovirus enhancement of transferrin-polylysine- mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8850- 8854. Liljistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semliki Forest Virus Replicon," (1991) Biotech. 9:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion- Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome- disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	183		/		
type 1 infection by cationic-liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. (1991) 72:2685-2696. Curiel, D.T. et al. "Adenovirus enhancement of transferrin-polylysine-mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8850-8854. Liljistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semliki Forest Virus Replicon," (1991) Biotech. 9:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	\$3			DNA into Cultured Cells with the Aid of Influenza Virus	
mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8830-8854. Liljistrom, P. and Garoff, H. "A New Generation of Animal Cell Expression Vectors Based on the Semliki Forest Virus Replicon," (1991) Biotech. 2:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	865		1	type 1 infection by cationic liposomes: the role of CD4, serum and	
Expression Vectors Based on the Semliki Forest Virus Repitcon, (1991) Biotech. 2:1356-1361. Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion- Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome- disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	X3		~	mediated gene delivery," (1991) Proc. Natl. Acad. Sci. USA 88:8850-	
Wirus," J. Cell Biology (1991) 112(4):615-623. Murata et al., "Modification of the N-Terminus of Membrane Fusion-Active Peptides Blocks the Fusion Activity," Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	86		<i>.</i> .	Expression Vectors Based on the Semliki Forest Virus Replicon, (1991)	
Active Peptides Blocks the Fusion Activity, "Biochem. and Biophys. Res. Communications (1991) 179(2):1050-1055. Cotton et al., (1992) "High-efficiency receptor-mediated delivery of small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles," Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	185			Phalen et al., "Cholesterol is Required for Infection by Semiki Forest Virus," J. Cell Biology (1991) 112(4):615-623.	
small and large 48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles, "Proc. Natl. Acad. Sci. USA 89:6094-6098. Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	SB		1	Active Peptides Blocks the Fusion Activity," Biochem. and Biophys.	
Adenovirus Coupled to DNA-Polylysine Complexes," Hum. Gene.	125		. /	small and large 48 kilobase gene constructs using the enaosome- disruption activity of defective or chemically inactivated adenovirus	
	Jac.			Curiel, D.T. et al., (1992) High-Efficiency Gene Transfer Mediated by Adenovirus Coupled to DNA-Polylysine Complexes, Hum. Gene. Therapy 3:147-154.	

EXAMINER

12 Busit

DATE CONSIDERED

*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

12/20/89

(A 100)	.*		Sheet 3 of 3			
FORM PTO MARK OFF.						
ATTY DOCKET NO. 32-95	SERIAL NO. 08/477,354	FILING DATE June 7, 1995				
APPLICANT Hawley-Nelson et al.	APPLICANT Hawley-Nelson et al. GROUP Unassigned					

	OTHER	KIUK A	RT (including Author, Title, Date, Pertinent Pages, etc.)
BB			Wagner, E. et al., (1992) "Coupling of adenovirus to transferrin- polylysine/DNA complexes greatly enhances receptor-mediated gene delivery and expression of transfected genes," Proc. Natl. Acad. Sci. USA 89:6099-6103.
A.P.			Wagner, E. et al., (1992) "Influenza virus hemagglutinin HA-2 N- terminal fusogenic peptides augment gene transfer by transferrin- polylysine-DNA complexes: Toward a synthetic virus-like gene-transfer vehicle," Proc. Natl. Acad. Sci. USA <u>89</u> :7934-7938.
B			Epand et al, "Peptide models for the membrane destabilizing actions of viral fusion proteins," Biopolymers 32:309 (1992).
J&G		1	Walker et al., "Cationic lipids direct a viral glycoprotein into the class I major histocompatibility complex antigen-presentation pathway," Proc. Natl. Acad. Sci. USA (1992) <u>89</u> :7915-7918.
\$63		\	Ciccarone et al., "Cationic Liposome-Mediated Transfection of Eukaryotic Cells: High Efficiency Nucleic Acid Delivery with Lipofectin, Lipofectace", and Lipofectamine" Reagents, "FASEB J., Abstracts, (1993) 7(7):A1131, Abstract No. 454.
DC.		_	Yoshimura et al. "Adenovirus-mediated Augmentation of Cell Transfection with Unmodified Plasmid Vectors," J. Biol. Chem. <u>268</u> :2300 (1993).
SAD		1	"Transfection Reagent," Genetic Engineering News (15 June 1993), p.12, column 4.
pB		/	Kamata, H. et al., "Amphiphilic peptides enhance the efficiency of liposome-mediated DNA transfection," Nucl. Acids Res. <u>22</u> (3):536-537 (1994).
BB		./	Remy et al., "Targeted gene transfer into hepatoma cells with lipopolyamine-condensed DNA particles presenting galactose ligands: A stage toward artificial viruses," Proc. Natl. Acad. Sci., USA <u>92</u> :1744-1748 (1995).

EXAMINER D. Gruss DATE CONSIDERED
4/24/96

*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

12/20/89